

On class preserving automorphisms of groups of order 32

Deepak Gumber and Hemant Kalra

School of Mathematics and Computer Applications

Thapar University, Patiala - 147 004, India

E-mail: dkgumber@yahoo.com, happykalra26@gmail.com

Abstract

We study class preserving automorphisms of groups of order thirty two and prove that only two groups have non-inner class preserving automorphisms.

2000 Mathematics Subject Classification: 20D15, 20D45.

Keywords: Class preserving automorphism, Central automorphism.

1 Introduction

Let G be a finite group and let p be a prime. For $x \in G$, let x^G denote the conjugacy class of x in G and let $\text{Aut}(G)$ denote the group of all automorphisms of G . An automorphism α of G is called a class preserving automorphism if $\alpha(x) \in x^G$ for all $x \in G$. Observe that every inner automorphism of G is a class preserving automorphism. The set $\text{Aut}_c(G)$ of all class preserving automorphisms is a normal subgroup of $\text{Aut}(G)$, and the set $\text{Inn}(G)$ of all inner automorphisms is a normal subgroup of $\text{Aut}_c(G)$. Let $\text{Out}_c(G)$ denote the group $\text{Aut}_c(G)/\text{Inn}(G)$. In 1911, Burnside [2, p. 463] posed the question: Does there exist a finite group G which has a non-inner class preserving automorphism? In other words, whether there exists a finite group G for which $\text{Out}_c(G) \neq 1$? In 1913, Burnside [3] himself gave an affirmative answer to this question. He constructed a group G of order p^6 , p odd, such that $\text{Out}_c(G) \neq 1$. After this more groups were constructed such that $\text{Out}_c(G) \neq 1$ (see [5], [7], [11], [16] and [17]) and many authors found the groups for which $\text{Out}_c(G) = 1$ (see [4], [10], [12], [13], [14]). It follows from [8] that for extra special p -groups all class preserving automorphisms are inner and from [9] it follows that $\text{Out}_c(G) = 1$ for all groups G of order p^4 . Recently Yadav [19] studied the groups of order p^5 , p an odd prime. He proved that if G and H are two finite non-abelian isoclinic groups, then $\text{Aut}_c(G) \cong \text{Aut}_c(H)$. He also proved that $\text{Out}_c(G) = 1$ for all the groups G except two isoclinic families. In the present paper we show that out of 51 groups of order 32, there are only two groups for which $\text{Out}_c(G) \neq 1$. A list of groups of order 32 is available from Sag and Wamsley's work [15]. We use Sag and Wamsley's list and adopt the same notations for the nomenclature

and presentations of the groups. However, we write the generators 1,2,3, and 4 used in [15] respectively as x, y, z , and w . All the results proved in section 3 are summarised in the following theorem.

Theorem A *For all groups G of order 32, except for the groups G_{44} and G_{45} , $\text{Out}_c(G) = 1$.*

By $\text{Hom}(G, A)$ we denote the group of all homomorphisms of G into an abelian group A . A non-abelian group G that has no non-trivial abelian direct factor is said to be purely non-abelian. For two subgroups H and K of G , $[H, K]$ denotes the subgroup of G generated by all commutators $[x, y] = x^{-1}y^{-1}xy$ with $x \in H$ and $y \in K$. By $[x, G]$ we denote the set of all commutators of the form $[x, g]$, $g \in G$. Observe that if G is nilpotent of class 2, then $[x, G]$ is a subgroup of G . The lower central series of a group G is the descending series $G = \gamma_1(G) \geq \gamma_2(G) \geq \dots \geq \gamma_i(G) \geq \dots$, where $\gamma_{n+1}(G) = [\gamma_n(G), G]$, and upper central series is the ascending series $Z(G) = Z_1(G) \leq Z_2(G) \leq \dots \leq Z_i(G) \leq \dots$, where $Z_{i+1}(G) = \{x \in G \mid [x, y] \in Z_i(G) \text{ for all } y \in G\}$. An automorphism α of G is called a central automorphism if $x^{-1}\alpha(x) \in Z(G)$ for each $x \in G$. The set $\text{Cent}(G)$ of all central automorphisms of G is a normal subgroup of $\text{Aut}(G)$.

2 Toolbox

All the tools required to prove Theorem A are listed in this section. Fortunately (or unfortunately!) we have been able to prove the theorem using the existing tools. Only one existing tool (Lemma 2.1) is slightly modified (Lemma 2.2).

Lemma 2.1 [19, Theorem 3.5] *Let G be a finite p -group of class 2 and let $\{x_1, x_2, \dots, x_m\}$ be a minimal generating set for G such that $[x_i, G]$ is cyclic, $1 \leq i \leq m$. Then $\text{Out}_c(G) = 1$.*

Lemma 2.2 *If G is a group of order p^n , $n \geq 3$ and $|Z(G)| = p^{n-2}$, then $\text{Out}_c(G) = 1$.*

Proof. Observe that nilpotency class of G is 2. If x is a non central element of G , then $|C_G(x)| = p^{n-1}$ and thus $|x^G| = |[x, G]| = p$. If x is a central element of G , then $|x^G| = |[x, G]| = 1$. In any case, $[x, G]$ is cyclic and therefore $\text{Out}_c(G) = 1$ by Lemma 2.1. $\square$

Lemma 2.3 [9, Proposition 4.1] *Let G be a group which can be generated by two elements x, y such that every element of G can be written in the form $x^i y^j$, i, j are integers. Then $\text{Out}_c(G) = 1$.*

Lemma 2.4 [8, Proposition 2.2] *Let $G = H \oplus K$ be the direct sum of its subgroups H and K . Then $\text{Out}_c(G) = 1$ if and only if $\text{Out}_c(H) = \text{Out}_c(K) = 1$.*

Lemma 2.5 [6, Proposition 2.7] *Let G be a finite group having an abelian normal subgroup A with cyclic quotient G/A . Then class-preserving automorphisms of G are inner automorphisms*

Lemma 2.6 [19, Lemma 2.2] *Let G be a finite p -group such that $Z(G) \subseteq [x, G]$ for all $x \in G - \gamma_2(G)$. Then $|\text{Aut}_c(G)| \geq |\text{Cent}(G)||G/Z_2(G)|$.*

Lemma 2.7 [1] *If G is a purely non abelian finite group, then $|\text{Cent}(G)| = |\text{Hom}(G/\gamma_2(G), Z(G))|$.*

The well known commutator identities

$$[x, yz] = [x, z][x, y][x, y, z]; \quad [xy, z] = [x, z][x, z, y][y, z],$$

where $x, y, z \in G$, will be frequently used without any reference.

3 Proof of Theorem A

The groups G_1 to G_7 are abelian, so we start with non abelian groups. The proof of the theorem follows from all the results proved in this section.

Proposition 3.1 *For groups G_{44} and G_{45} , there exist class preserving automorphisms which are not inner.*

Proof. Consider $G = G_{44}$. Since $yx = xy^5$ and $zy = y^7z$, it follows that $|y| = 8$ and every element of G can be written in the form $x^i y^j z^k$, where $0 \leq i, k \leq 1$, and $0 \leq j \leq 7$. We show that $Z(G) = \{1, y^4\}$. Since $[z, y^4] = [z, y^2]^2 = [z, y]^4 = y^8 = 1$ and $[x, y^2] = [x, y]^2 = y^8 = 1$, $y^4 \in Z(G)$. It is easy to see that other powers of y are not in $Z(G)$. Since xy^4 does not commute with y , it cannot be in $Z(G)$. If $xy^i \in Z(G)$; $i = 1, 2, 3, 5, 6, 7$, then commuting it with z gives $i \equiv 7i \pmod{8}$, a contradiction. Using similar arguments, other possibilities can also be ruled out and thus $Z(G) = \{1, y^4\}$. Consider $G/Z(G)$. Since $xZ(G)$ commutes with $yZ(G)$ and $zZ(G)$, and $y^2Z(G)$ commutes with $xZ(G)$ and $zZ(G)$, it follows that $|Z(G/Z(G))| \geq 4$. Since $yZ(G) \notin Z(G/Z(G))$, $|C_{G/Z(G)}yZ(G)| = 8$ and thus $|Z(G/Z(G))| = 4$. This gives $|Z_2(G)| = 8$. Now $G/Z_2(G)$ is abelian, therefore $G = Z_3(G)$ and hence G is nilpotent of class 3. It is easy to see that $Z(G) \leq \gamma_2(G) = \Phi(G)$. It follows from [18, theorem 4.7] that $|\gamma_2(G)| = 4$ and $Z(G) \subseteq [x, G]$ for all $x \in G - \gamma_2(G)$. By Lemma 2.6

$$|\text{Aut}_c(G)| \geq |\text{Cent}(G)|2^5/|Z_2(G)|.$$

Since G is purely non abelian and $G/\gamma_2(G)$ is elementary abelian of order 8, by Lemma 2.7 we have

$$|\text{Cent}(G)| = |\text{Hom}(G/\gamma_2(G), Z(G))| = 8.$$

Thus $|\text{Aut}_c(G)| \geq 2^5 > 2^4 = |\text{Inn}(G)|$. That $\text{Out}_c(G_{45}) \neq 1$ can be shown similarly. $\square$

Remark: The order of $\text{Aut}_c(G)$ in the above proposition is in fact exactly equal to 32. Observe that $C_G(x) = \langle x, y^2, z \rangle$, $C_G(y) = \langle y \rangle$ and $C_G(z) = \langle x, y^4, z \rangle$. Thus $|C_G(x)| = 16$ and $|C_G(y)| = |C_G(z)| = 8$. Hence $|x^G| = 2$

and $|y^G| = |z^G| = 4$. Since any class preserving automorphism preserves the conjugacy classes, there are $|x^G|$, $|y^G|$ and $|z^G|$ choices for the images of x , y and z respectively under it. Thus $|\text{Aut}_c(G)| \leq |x^G||y^G||z^G| = 32$.

Proposition 3.2 *The groups G_{42} and G_{43} are extraspecial and therefore all class preserving automorphisms are inner.*

Proof. Consider $G = G_{42}$. Since $1 = [y^2, w] = [x^2, w] = [x, w]^2 = x^4$, it follows that $|x| = |y| = |z| = |w| = 4$ and $x^2 \in Z(G)$. Since $wx = xw^3$ and $zy = y^3z$, every element of G can be written as $x^i y^j z^k w^l$, where $0 \leq i \leq 3$ and $0 \leq j, k, l \leq 1$. If $z(x^i y) = (x^i y)z$, then $x^i y^3 z = x^i y z$ and thus $y^2 = 1$ which is not so. So $x^i y \notin Z(G)$. Using similar arguments we can show that $x^i z, x^i w, x^i y z, x^i y w, x^i z w$, and $x^i y z w$ cannot be in the center. Thus $Z(G) = \{x^2, 1\}$. Observe that any element of $G/Z(G)$ is of the form $x^i y^j z^k w^l Z(G)$, where $0 \leq i, j, k, l \leq 1$. Since $xw = wx^3 \equiv wx \pmod{Z(G)}$ and $yz = zy^3 \equiv zy \pmod{Z(G)}$, it follows that $G/Z(G)$ is abelian and hence $\gamma_2(G) = Z(G) = \Phi(G)$. Similar arguments show that G_{43} is also an extraspecial p -group. $\square$

Proposition 3.3 *If G is any of the groups from G_{23} to G_{25} , then $G = H \oplus K$ for some subgroups H and K such that $\text{Out}_c(H) = \text{Out}_c(K) = 1$ and hence $\text{Out}_c(G) = 1$ by Lemma 2.4.*

Proof. Observe that $G_{23} = H \oplus K$, where $H = \{x, y | x^8 = y^2 = 1, yx = x^7 y\}$ and $K = \{z | z^2 = 1\}$; $G_{24} = H \oplus K$, where $H = \{x, y | x^8 = y^2 = 1, yx = x^3 y\}$ and $K = \{z | z^2 = 1\}$; and $G_{25} = H \oplus K$, where $H = \{x, y | x^8 = 1, y^2 = x^4, y^{-1}xy = x^{-1}\}$ and $K = \{z | z^2 = 1\}$. Since K is cyclic and every element of H can be written as $x^i y^j$ for suitable i and j , $\text{Out}_c(K) = \text{Out}_c(H) = 1$. $\square$

Proposition 3.4 *If G is any of the groups from G_{22} , G_{29} , G_{30} , G_{32} , and $G_{49} - G_{51}$, then every element of G can be written as $x^i y^j$, for suitable integers i and j , and hence $\text{Out}_c(G) = 1$ by Lemma 2.3.*

Proof: Observe that all the groups are generated by x and y , and yx can be written as $xy^9, x^7 y, x^3 y, x^7 y, x^{15} y, x^7 y$ and $x^{15} y$ in the respective groups. $\square$

Proposition 3.5 *If G is any of the groups from G_8 to G_{21} , then $|Z(G)| = 8$ and hence $\text{Out}_c(G) = 1$ by Lemma 2.2.*

Proof. For G_8 it is clear that z and w are in the center. Since $[x^2, y] = [x, y]^2 = 1$, x^2 is in the center and thus $|Z(G_8)| = 8$. For G_9 again it is obvious that z and w are in the center. Since $x^4 = [y, x]^2 = [y^2, x] = 1$, $|x| = 4$ and since $[x^2, y] = [x, y]^2 = 1$, x^2 is in the center and thus $|Z(G_9)| = 8$. For G_{10} , x and w are in the center and since $x^4 = [z, y]^2 = [z^2, y] = 1$, $|x| = 4$ and therefore $|Z(G_{10})| = 8$. For G_{11} , clearly z is in the center and since $[x, y^2] = [x, y]^2 = [x^2, y] = 1$, it follows that y^2 is in the center and $|[x, y]| = 2$. Also, since z commutes with x and y , it commutes with $[x, y]$ as well. Thus $[x, y]$ is also in the center and hence $|Z(G_{11})| = 8$. For G_{12} , z is in the center

and since $[x^2, y] = [x, y^2] = 1$, both x^2 and y^2 are in the center and thus $|Z(G_{12})| = 8$. Observe that every element of G_{13} can be written in the form $x^i y^j z^k$; $0 \leq i \leq 1$, $0 \leq j \leq 7$, and $0 \leq k \leq 1$. Therefore $|y| = 8$ and $|x| = 2$. Now clearly z is in the center and $[y^2, x] = [y, x]^2 = 1$ implies that y^2 is also in the center. Therefore $|Z(G_{13})| = 8$. For G_{14} , clearly z is in the center and since $[x^2, y] = [x, y]^2 = 1$, x^2 is in the center and thus $|Z(G_{14})| = 8$. In G_{15} , observe that $x^4 = [y, x]^2 = [y^2, x] = 1 = [y, x^2]$. Thus $|x| = 4$ and x^2 is in the center. Also z is in the center and therefore $|Z(G_{15})| = 8$. In G_{16} , $|x| = 4$ and x and z^2 are in the center. Thus $|Z(G_{16})| = 8$. In G_{17} , x is in the center and $|x| = 8$. Thus $|Z(G_{17})| = 8$. In G_{18} , $[x, y]$ is in the center. Also $[x^2, y] = [x, y]^2 = [x, y]^2 = 1$ implies that x^2 and y^2 are in the center and $|[x, y]| = 2$. Therefore $|Z(G_{18})| = 8$. In G_{19} , since $[x^2, y] = [x, y]^2 = [x, y]^2 = 1$, it follows that x^2 and y^2 are in the center and $|Z(G_{19})| = 8$. In G_{20} clearly $[x, y]$ is in the center and $[x, y^2] = [x, y]^2 = [x^2, y] = 1$ implies that $|[x, y]| = 2$ and y^2 is in the center. Thus $|Z(G_{20})| = 8$. In G_{21} it follows from the relation $[x^2, y] = [x, y^2] = 1$ that x^2 and y^2 are in the center and $|Z(G_{21})| = 8$. $\square$

Proposition 3.6 *If G is any of the groups from $G_{26} - G_{28}$, G_{31} , $G_{33} - G_{41}$ and $G_{46} - G_{48}$, then G has an abelian normal subgroup H such that G/H is cyclic and hence $\text{Out}_c(G) = 1$ by Lemma 2.5.*

Proof. In G_{26} , $1 = [x^2, z] = [y^4, z] = [y^2, z]^2 = [y, z]^4$. Thus $|x| = 4$, $|y| = 8$, and $H = \langle x, y \rangle$. In G_{27} , $1 = [x, y^2] = [x, y]^4 = x^8$ implies that $|x| = 8$. Let $z = [x, y]$, then $[z, x] = 1$ and $|z| = 4$. Thus $H = \langle x, z \rangle$. In G_{28} , $1 = [x^4, y] = [x^2, y]^2 = [x, y]^4 = x^8$ implies that $|x| = 8$. Take $z = [x, y]$, then $[z, x] = 1$, $|z| = 4$ and $H = \langle x, z \rangle$. In G_{31} , take $z = [x, y]$, then z commutes with x . Also, $1 = [x^4, y] = [x^2, y]^2 = [x, y]^4 = z^4$ implies that $|z| = 4$ and thus $H = \langle x, z \rangle$. In G_{33} , observe that each of the elements x, y, zxz^{-1} , and zyz^{-1} is of order 2 and commutes with another. Thus $H = \langle x, y, zxz^{-1}, zyz^{-1} \rangle$. In G_{34} , it is easily seen that $H = \langle x, y \rangle$. In G_{35} , observe that $|x| = 4$ and $H = \langle x, y \rangle$. In G_{36} and G_{37} , $1 = [x, z]^2 = [x^2, z]$ and $[x, z] = x^2$ implies that $xz = zx^3$. Take $w = [y, z]$, then $|w| = 2$ and it commutes with x and y . Thus $H = \langle x, y, w \rangle$. In G_{38} , $1 = [x^2, z] = [x, z]^2 = y^4$; $(y^2 z)^2 = ([z, x] z)^2 = 1$. Thus $|y| = 4$ and $y^2 z = zy^2$. Take $w = [y, z]$, then $|w| = 2$ and it commutes with y and z . Thus $H = \langle x, y, w \rangle$. In G_{39} , $[x^2, z] = [x, z]^2 = x^4$; $1 = [x, z^2] = [x, z]^2 [x^2, z] = x^8$; $[y^2, z] = [y, z]^2 = x^4$; $1 = [y, z^2] = [y, z]^2 [y, z, z] = x^4 [x^2 y^2, z] = x^4 [x^2, z] [y^2, z] = x^{12}$. Thus $|x| = 4$ and $H = \langle x, y \rangle$. In G_{40} , $z^4 = [z, y]^2 = [z^2, y] = [x^{-2} y^2, y] = 1$; $[y^2, z] = [y, z]^2 = 1$; $x^4 = [x, z]^2 = [x^2, z] = [y^2 z^2, z] = 1$. Thus $|x| = 4$, $|y| = 4$ and $H = \langle x, y \rangle$. In G_{41} , $[x^2, z] = [x, z]^2 = x^4 y^4$; $[y^2, z] = [y, z]^2 = x^4$; $1 = [x, z^2] = [x, z]^2 [x^2 y^2, z] = x^{12} y^8$; $1 = [y, z^2] = [y, z]^2 [x^2, z] = x^8 y^4$. Thus $|x| = 4$, $|y| = 4$ and $H = \langle x, y \rangle$. In G_{46} , let $z = [x, y]$ and $w = [x, y, y] = [z, y]$, then z commutes with x and w commutes with y . Observe that $1 = [x^2, y] = [x, y]^2 = z^2$; $[x, y^2] = [x, y]^2 [x, y, y] = w$; $1 = [x, y^4] = [x, y^2]^2 [x, y^2, y^2] = w^2$. Thus $|z| = |w| = 2$. Since $xy^2 = (xy)y = (yxz)y = yx(zy) = yx(yzw) = y(xy)zw = y(yxz)zw = y^2 xw$, $wx = (xy^2 xy^2)x = (xy^2 x)y^2 x = (xy^2 x)(xwy^2) = xw$. Thus w commutes

with x and hence with z as well. It is easy to see that every element of G_{46} can be written in the form $x^i y^j z^k w^l$, where $0 \leq i, k, l \leq 1$ and $0 \leq j \leq 3$. Let $H = \langle x, z, w \rangle$, then H is abelian, $Hy = yH$ and $G_{46}/H = \langle yH \rangle$. In G_{47} let $[x, y] = z$, then z commutes with x and $[z, y] = y^4$. Observe that $1 = [x^2, y] = [x, y]^2 = z^2$; $[z, y^2] = [z, y]^2 = y^8$; and $1 = [z^2, y] = [z, y][z, y, z][z, y] = y^4[y^4, z]y^4 = y^4[y^2, z]^2y^4 = y^{-8}$. Thus y^2 commutes with z , $|z| = 2$ and $|y| = 8$. Since $zy = y^5z$ and $yx = xyz$, every element of G_{47} can be written as $x^i y^j z^k$, where $0 \leq i, k \leq 1$ and $0 \leq j \leq 7$. Let $H = \langle x, y^4, z \rangle$, then it is easy to verify that H is a normal subgroup of G_{47} and $G_{47}/H = \langle Hy \rangle$ is cyclic. In G_{48} , let $[x, y] = z$, then z commutes with x and $z^2 = [x^2, y] = 1$. Since $1 = [z^2, y] = [z, y]^2 = y^8$, $|x| = 4$ and $|y| = 8$. Now $zy = y^5z$ and $yx = xyz$, therefore every element of G_{48} can be written as $x^i y^j z^k$, where $0 \leq i, k \leq 1$ and $0 \leq j \leq 7$. Let $H = \langle x, z \rangle$, then $Hy = yH$ and $G_{48}/H = \langle Hy \rangle$. $\square$

Acknowledgement: The research of the second author is supported by Council of Scientific and Industrial Research, Government of India. The same is gratefully acknowledged.

References

- [1] J. E. Adney and T. Yen, *Automorphisms of a p -group* Illinois J. Math. **9** (1965), 137-143.
- [2] W. Burnside, *Theory of groups of finite order*, 2nd Ed. Dover Publication, Inc., 1955. Reprint of the 2nd edition (Cambridge, 1911).
- [3] W. Burnside, *On the outer automorphisms of a group*, Proc. London Math. Soc. (2) **11** (1913), 40-42.
- [4] M. Fuma and Y. Ninomiya, “*Hasse principle*” for finite p -groups with cyclic subgroups of index p^2 , Math. J. Okayama Univ. **46** (2004), 31-38.
- [5] H. Heineken, *Nilpotente Gruppen, deren sämtliche Normalteiler charakteristisch sind*, Arch. Math. (Basel) **33** (1980), No. 6, 497-503.
- [6] M. Hertweck and E. Jespers, *Class-preserving automorphisms and normalizer property for Blackburn groups*, J. Group Theory **12** (2009), 157-169.
- [7] D. Jonah and M. Konvisser, *Some non-abelian p -groups with abelian automorphism groups*, Arch. Math. (Basel) **26** (1975), 131-133.
- [8] M. Kumar and L. R. Vermani, “*Hasse principle*” for extraspecial p -groups, Proc. Japan Acad. **76**, Ser. A, No.8 (2000), 123-125.
- [9] M. Kumar and L. R. Vermani, “*Hasse principle*” for the groups of order p^4 , Proc. Japan Acad. **77**, Ser. A, No. 6 (2001), 95-98.
- [10] M. Kumar and L. R. Vermani, *On automorphisms of some p -groups*, Proc. Japan Acad. **78**, Ser. A, No. 4 (2002), 46-50.

- [11] I. Malinowska, *On quasi-inner automorphisms of a finite p -group*, Publ. Math. Debrecen **41** (1992), No. 1-2, 73-77.
- [12] T. Ono, *“Hasse principle” for $GL_2(D)$* , Proc. Japan Acad. **75** Ser. A (1999), 141-142.
- [13] T. Ono and H. Wada, *“Hasse principle” for free groups*, Proc. Japan Acad. **75** Ser. A (1999), 1-2.
- [14] T. Ono and H. Wada, *“Hasse principle for symmetric and alternating groups*, Proc. Japan Acad. **75** Ser. A (1999), 61-62.
- [15] T. W. Sag and J. W. Wamsley, *Minimal presentations for groups of order 2^n , $n \leq 6$* , J. Aust. Math. Soc. **15** (1973), 461-469.
- [16] C. H. Sah, *Automorphisms of finite groups*, Journal of Algebra **10** (1968), 47-68.
- [17] F. Szechtman, *n -inner automorphisms of finite groups*, Proc. Amer. Math. Soc. **131** (2003), 3657-3664.
- [18] H. Tendra and W. Moran, *Flatness conditions on finite p -groups*, Comm. Algebra **32** (2004), 2215-2224.
- [19] M. K. Yadav, *On automorphisms of some finite p -groups*, Proc. Indian Acad. Sci. (Math Sci.) **118** (1) (2008), 1-11.